

## Review article

# Schistosomosis in Burkina Faso: review on 60 years' research

Adama ZIDA<sup>1,2,3</sup>, Thierry K. GUIGUEMDE<sup>2,4</sup>, Marcel P. SAWADOGO<sup>1,2</sup>,  
Chanolle TCHEKOUNOU<sup>5</sup>, Issiaka SOULAMA<sup>3,6</sup>, Cheick S. BARRY<sup>2,5</sup>,  
Mohamadou SIRIBIE<sup>7</sup>, Florian MARKS<sup>7,8,9</sup>, Ibrahim SANGARÉ<sup>10</sup>,  
Bamba SANATA<sup>10</sup>

<sup>1</sup>Service de Parasitologie-Mycologie, Centre Hospitalier Universitaire Yalgado Ouedraogo, Ouagadougou, Burkina Faso

<sup>2</sup>Ecole Doctorale Sciences et Santé, Université Joseph Ki-Zerbo, Burkina Faso

<sup>3</sup>Centre National de Recherche et de Formation sur le Paludisme, Ouagadougou, Burkina Faso

<sup>4</sup>Centre Hospitalier Universitaire Charles De Gaulles, Ouagadougou, Burkina Faso

<sup>5</sup>Institut International des Sciences et Technologie (IISTech), Ouagadougou, Burkina Faso

<sup>6</sup>Institut de Recherche en Sciences de la Santé (IRSS/CNRST), Ouagadougou, Burkina Faso

<sup>7</sup>International Vaccine Institute (IVI), Seoul, Republic of Korea

<sup>8</sup>Cambridge Institute of Therapeutic Immunology and Infectious Disease, University of Cambridge School of Clinical Medicine, Cambridge, UK

<sup>9</sup>University of Antananarivo, Antananarivo, Madagascar

<sup>10</sup>Institut Supérieur des Sciences de la Santé, Université Nazi Boni, Burkina Faso

Corresponding Author: Adama Zida; e-mail: zidaadama@live.fr

**ABSTRACT.** Schistosomosis is a common neglected helminthic disease in the tropics and sub-tropics particularly in sub-Saharan countries including Burkina Faso. It is the second world parasitic endemic disease after malaria. The two prevalent species infecting human in Burkina Faso are *Schistosoma haematobium* and *Schistosoma mansoni* which cause respectively the urogenital schistosomosis and the intestinal schistosomosis. This review aimed at providing an historical perspective of research on schistosomosis from 1960 to 2020 and shedding some light on the gaps in knowledge useful for the disease control and the elimination efforts in Burkina Faso. Formal systematic review was not followed for this review. Published studies on the schistosomosis in Burkina Faso over the period from 1960 to 2020, were search in Medline, PubMed, Google Scholar, EMBASE and the libraries of main universities in Burkina Faso namely: Joseph Ki-Zerbo University and Nazi Boni University. The following keywords used were: schistosomiasis, schistosomosis, *Bilharzia*, *Bulinus*, *Biomphalaria*, Upper-Volta and Burkina Faso. Over a period of 60 years, a total of 87 scientific research documents were identified. The original scientific research articles represent the majority of the scientific documents found (65.52%). Urinary schistosomosis was the most common from the documentation. There has been a gradual decrease in the prevalence, more significantly since the implementation of the National Schistosomiasis Control Program (NSCP). The effectiveness of the NSCP could therefore contribute to the elimination of schistosomosis in Burkina Faso.

**Keywords:** schistosomosis, bilharziosis, *Schistosoma*, *Bulinus*, *Biomphalaria*, Burkina Faso

## Introduction

Schistosomosis or bilharzia is one of the most widespread parasitic diseases in the world. Schistosomosis is the second world parasitic endemic after malaria. Nearly 800 million people

around the world are at risk of infection [1]. Schistosomes are responsible for 800,000 deaths per year worldwide. In 52 countries of the world, 230 million people need annual treatment, i.e., 80 to 90% live in Africa [1]. In 2003, in sub-Saharan Africa, 70 million individuals with hematuria and

32 million of those with dysuria were infected with *Schistosoma haematobium*. *Schistosoma mansoni* is believed to be the cause of diarrhea in 0.78 million people, melena in 4.4 million cases and hepatomegaly in 8.5 million cases [2]. These schistosomes are endemic in 76 countries, most of which are located in Africa. Thus, urinary schistosomiasis caused by *S. haematobium* affects 54 countries in Africa and the eastern Mediterranean basin [3]. WHO stated in 2006 that some 200 million people are infected with schistosomiasis in 76 developing countries around the world and that about 80% of cases come from Africa [4]. In Burkina Faso, in the years 1997–1998, the overall prevalence of urinary bilharzia was about 30% [5]. Concerning intestinal bilharzia, its prevalence varied between 36–79% in the 1980s [6].

Taking into account the creation of water reservoirs in the hydraulic installations development (large dams, irrigation networks, small community dams) for agricultural and or energy needs, and the increase in anthropogenic pressure, man creates ecological disturbances that can lead to the proliferation of initially rare biological species; in particular the intermediate host molluscs of schistosomes [5,7,8]. For this purpose, we can estimate the population reached by all forms of bilharzia to be 3 million people in Burkina Faso [9]. Although, the lethality associated with schistosomiasis is low in Burkina Faso (rate of 0.34%) [10], the extent of this disease is measured particularly in developing countries where poverty evolves in parallel with malnutrition, lack of safe water and precarious health infrastructure [11–13]. Diseases by extension, the importance of bilharzia is however underestimated in public health: their distribution in the population is very heterogeneous and they remain clinically asymptomatic or discrete for a long time, limited to hematuria for the urinary form or diarrhea for the intestinal form. Many works have been carried out by authors on schistosomiasis but still under exploited. The aim of this work is therefore to contribute to the popularization of the results of research on schistosomiasis in Burkina Faso from January 1960 to December 2020.

## Methods

Our present review study is concerned only with schistosomiasis in Burkina Faso; a landlocked country located in the heart of West Africa. It covers an area of approximately 274,200 km<sup>2</sup> and is

located between latitudes 9°20' and 15°05' North and longitudes 5°30' West and 2°20' East. Burkina Faso, is a Sahelian country bounded to the Est by Niger, to the North and West by Mali, to the South by Ghana, Côte d'Ivoire, Togo and Benin. The main rivers of the country are the Mouhoun river, the Nakambe river, the Nazinon river and the Comoe river. The rivers are connected to three main basins: the Volta, Comoe and Niger basins. The vast majority of the Burkinabe population, about 77.3%, lives in rural areas with agriculture as their main activity [14].

This is a retrospective study based on the review of scientific documents that addressed schistosomiasis in Burkina Faso from January 1960 to December 2020. The review took place from January to December 2021. Our study included any scientific document related to schistosomes and their intermediate hosts in Burkina Faso published online or available in libraries from January 1960 to December 2020. We concluded the study on 31st December 2021. Finally, we considered the studies eligible only if they met the following inclusion criteria: a) schistosomiasis and other pathologies in Burkina Faso, b) schistosomiasis both Burkina Faso and other countries.

Our data has been collected using the search database such as: Medline, Google Scholar, PubMed, EMBASE and the libraries of our various universities namely: Joseph Ki-Zerbo University and Nazi Boni University. The key words used were: schistosomiasis, schistosomiasis, bilharzia, *Bulinus*, *Biomphalaria*, Upper-Volta and Burkina Faso.

The informations were collected using individual collection sheet for each listed scientific document. The following variables were considered: a) quantitative variables: the prevalence and year of publication, b) qualitative variables: title of the documents, the type of the document, the nationality of the authors, the sources of funding, the species of schistosomes and their intermediate hosts, the regions, the provinces and the study sites, the populations of the study, the parasitological, immunological and anatomopathological techniques carried out.

The prevalence rates were calculated using SPSS version 17 software. The histograms and pie charts were generated using Excel 2010 software.

## Results

### *Inventory of scientific documents on schistosomosis according to their nature from January 1960 to December 2020*

Over a period of 60 years (1960 to 2020), we have identified 87 scientific documents on schistosomosis that meet the inclusion criteria. The nature of the documents listed was dominated by original scientific research articles (65.51%) (Fig. 1).




Figure 1. Distribution of scientific documents on schistosomosis according to their nature from January 1960 to December 2020

### *Number of documents listed by decade*

We have noted an increase in the number of scientific documents addressing schistosomosis over the time from 1960 to 2020 (Tab. 1).

Table 1. Number of documents listed by decade

Period	Number (n=87)	Proportion (%)	References
1960–1969	01	1.15	[15]
1970–1979	05	5.75	[16–20]
1980–1989	14	16.09	[21–34]
1990–1999	16	18.39	[35–50]
2000–2009	24	27.59	[51–74]
2010–2020	27	31.03	[75–101]

### *Distribution of documents according to the subject of the study and the study population*

Despite the multiplicity of intermediate hosts, the majority of the studies (60.92%) conducted were focused on schistosomes. It is to be noted that 29.89% of the documents addressed schistosomes and intermediate hosts, while very few scientific documents (4.60%) were interested in studying the behavior of intermediate hosts (Tab. 2).

During the period covered by the study, 49.40% of the scientific documents mentioned surveys were carried out on the population in general regardless of age or gender.

Table 2. Distribution of documents according to the study population and the study subject

Biologic examination	Number of documents	Proportion (%)	References
Study population	83	100	
Population in general	41	49.40	[1–4,6,8,9,12,15–18,20–23,25–27,29,35,37,38,41,47,49,51,54, 56, 58,59,64,69,78,83,85,86]
Children	35	42.17	[19,29,30,31,34,35,37,38,41,43, 47,49,51,54,56, 58,59,64,69,72, 76,77,79,81,82,83, 84,87,89,91,93,99,103]
Women	04	4.82	[74–76,102]
Traveller	03	3.61	[48,58,62]
Study subject	87	100	
Schistosomes	55	63.21	[1–4,6,9,11,12,15–19,21,23, 25,27,29,30,31, 34,35,37,38,43, 47–54,56–59, 63,65,66,67,74,78,79, 82,84, 87,91,93,96,97,98,101, 102,103]
Schistosomes and intermediate hosts	27	31.03	[26,41,42,45,55, 60,68,69,75,76,77,80,81,83,85,86,88,89, 90,92, 94,95,97,99,100]
Intermediate hosts	5	5.74	[20,22,40,44,46]

### Distribution of documents according to schistosome species

Three species of the genus *Schistosoma*, namely *S. mansoni*, *S. haematobium* and *S. intercalatum*, all parasites of humans, were highlighted in studies carried out in Burkina Faso from 1960 to 2020. The documents concerning both *S. mansoni* and *S. haematobium* had the highest percentage (50.60%). Individually, *S. haematobium* was the most studied species with a percentage of 37.35. This distribution is illustrated in Figure 2.




Figure 2. Distribution of documents according to the species of schistosomes

### Distribution of documents according to biological and anatomopathological diagnostic techniques

The urinary filtration and Kato-Katz techniques were more associated in scientific documents with

Table 3. Distribution of documents according to biological and anatomopathological diagnostic techniques

Biologic examination	Number	Proportion (%)	References
Urinary filtration			
and Kato-Katz	40	48.19	[9,16,19,23,25,27,29,37,38,41,43,47,48,49,51,54,55,57,58,59,64,67,68,74,75,77,81,83,85,86,89,90–95,97–99]
Urinary filtration	25	30.12	[18,21,28,30,31,34–36,50,52,53,60,63,66,69,73,76,79,80,83,84,96,100–102]
Kato-Katz	07	8.43	[15,17,26,56,63,78,82]
Anatomopathology	04	4.82	[32,50,72,87]
Indirect hemagglutination	02	2.41	[12,71]
Indirect immunofluorescence	02	2.41	[24,71]
Immunohistochemistry/PCR	02	2.41	[12,65]
ELISA test	01	1.21	[65]
Total	83	100	

48.19%. Individually, the urinary filtration technique was the most diagnostic method applied with 30.12% (Tab. 3).

### Distribution of documents according to intermediate hosts found by region

The following species: *Bulinus truncatus*, *B. senegalensis*, were found in 13 regions of Burkina Faso, including respectively 12 and 11 documents that mentioned the Mouhoun Loop. *Biomphalaria pfeifferi* was found sporadically in 12 regions of Burkina Faso (Tab. 4).

### Distribution of documents according to the molecules evaluated

Out of the total of 83 documents, we observed only 6 (6.89%) that mentioned evaluated molecules. Praziquantel was the most experimented molecule with 3 out of the 6 documents listed (50%), followed by oxamniquine with 2 documents (33.33%) and metrifonate with one document (16.67%).

### Distribution of documents according to the state and funding source

Only 13 studies out of the 87 noticed (14.94%) had been funded. Regarding the funding sources, the AUF and the USAID are the first founding institutions with 5 investigations funded each (38.46%) (Tab. 5).

Table 4. Distribution of documents according to intermediate hosts found by region

Intermediate hosts	Regions and number of documents												
	Hauts- bassins	Cascades	Sud-ouest	Boucle du Mouhoun	Centre	Plateau central	Centre-nord	Centre-sud	Centre-ouest	Centre-est	Est	Nord	Sahel
	n	n	n	n	n	n	n	n	n	n	n	n	n
<i>Bulinus truncatus</i>	2	3	1	12	3	4	4	1	2	2	3	1	1
<i>B. senegalensis</i>	2	2	1	11	3	2	2	1	2	1	2	1	1
<i>B. globosus</i>	3	1	1	–	2	1	2	1	1	2	2	1	1
<i>B. forskalii</i>	1	1	1	4	1	1	1	1	1	1	1	1	–
<i>B. umbilicatus</i>	–	–	–	–	1	1	1	1	1	1	2	–	–
<i>B. genius</i>	1	–	–	–	–	–	–	–	–	–	–	–	–
<i>B. tropicus</i>	–	–	–	–	–	–	1	–	–	–	–	–	–
<i>B. jousseaumei</i>	1	–	–	–	–	–	–	–	–	–	–	–	–
<i>B. pfeifferi</i>	3	2	1	7	1	2	1	1	1	1	2	–	1

Table 5. Distribution of documents according to their financing status

	Number of documents	Proportion (%)
Status	87	100
Mention of financing	13	14.94
No mention of financing	74	85.05
No mention of financing		
Funding sources	13	100
AUF	05	38.46
USAID	05	38.46
DBL	02	15.38
APARET	01	7.70

#### *Average prevalence of schistosomosis in Burkina Faso by decade according to the documents used*

AUF: Agence Universitaire de la Francophonie; USAID: United States Agency for International Development; DBL: Danish Bilharziasis Laboratory-Institute for Health Research

and Development; APARET: African Program for Advanced and Research Epidemiology Training

We have observed a gradual decrease in the prevalence of schistosomosis over the years in Burkina Faso. *S. haematobium* and *S. mansoni*, being the two most frequently found species, each showed a clearly declining prevalence after these 60 years (Fig. 3).

#### Discussion

Our study is a literature review on schistosomosis in Burkina Faso from the year of independence (1960) to 2020, i.e. 60 years of existence. Eighty-seven (87) documents on schistosomosis have been identified during these 60 years in Burkina Faso. Our values are close to those of Makaula et al. [102] in Malawi in 2014 who identified 89 documents on schistosomosis over a period of 31 years. The establishment of the National Schistosomiasis Control Program (NSCP) in 2002 [97], the increase in the number of skills (doctoral students, researchers) and the sponsors' interests to the schistosomosis (NGOs, Research




Figure 3. Evolution of the average prevalence of schistosomiasis and *Schistosoma* species in Burkina Faso by decade

institutes) would have contributed to obtaining this documentary plurality on schistosomiasis in Burkina Faso. Original scientific research articles were the most frequently encountered type of document as evidenced by our results (65.52%). This could be explained by the easier access route (Internet) of scientific research results associated with their frequency of production compared to other types of documents observed.

Schistosomes were more studied than their intermediate hosts. This could be explained by the relative simplicity of the interventions on the definitive host (human) than on the intermediate host. In addition, this work on schistosomes in general has made it possible to identify the corresponding species and clinical forms. It allowed the scientific community to determine the prevalence and evolution of this neglected disease around water points. It also contributed to knowing and monitoring the level of schistosomiasis endemicity in the different regions of Burkina Faso. Most of the studies in our documents focused on the population in general (49.40%). The studies that were carried out on the population in general aimed to identify population groups at risk in order to better guide interventions from the national program. Indeed, Ouedraogo [34] in 1984, had demonstrated that in Kaya, Burkina Faso, the

schistosomiasis contamination risk was high for populations who practiced water collection, sale of brickyard water, livestock watering, market gardening, rice cultivation, maintenance of canals. And that, this risk was higher in those who were involved in swimming and fishing games. The studies conducted on children were not so negligible (42.17%). This is because children would be the most vulnerable population group.

The majority of the work (50.60%) concerned both urinary and intestinal schistosomiasis. This would be justified by the researchers' interest in comparing at the same time the prevalence of the two predominant forms of schistosomiasis on sentinel sites, but also in studying the prevalence of bi parasitism linked to the two species of schistosomes that are prevalent in Burkina Faso. It has been observed that urinary schistosomiasis has received much more attention from the scientific community in comparison with intestinal schistosomiasis. Indeed, 37.35% of the documents had to exclusively deal with urinary schistosomiasis against 10.84% of the documents for intestinal schistosomiasis. Then, it can be anticipated that urogenital schistosomiasis still remains a serious health problem in certain regions of Burkina Faso [101]. These results are similar to those of Makaula et al. [102] who found in their study more

documents concerning urinary schistosomosis than intestinal schistosomosis.

Urinary filtration as a diagnostic technique was the most used in the first documents found, followed by the Kato-Katz technique. Indeed, these techniques remained essential for researchers because they would make it possible to better conduct mass surveys and obtain good results. This idea is supported by the WHO which recommends the use of methods with lower sensitivity but very fast and less expensive [3].

The most listed intermediate hosts of *S. haematobium* and *S. mansoni* are the genus *Bulinus* and *Biomphalaria*, respectively. *Bulinus truncatus*, *B. globosus* and *B. senegalensis* have been identified in all ecological zones. *Biomphalaria pfeifferi* are found preferentially in the southern half of the country, which, with the exception of the Sourou hydro-developments, confirms the observation that the 14th northern parallel would be the limit of northern extension of these two species [6]. *B. umblicatus* found in 1994 in a few regions (Centre, Plateau central, Centre-Nord, Centre-Sud, Centre-Ouest, Centre-Est, Est) is inconstant because it is no longer found in the studies that followed [102]. *B. jousseaumei* and *B. genius*, found in the Hauts-bassins region respectively in 1987 and in 2012, *B. tropicus* found in the *Centre-Nord* region in 1992 are intermediate hosts of *S. haematobium* but very rarely found in Burkina Faso. Moreover, they were no longer found in the studies that followed [6].

According to the molecules tested, only six studies (6.89%) evaluated the molecules used in the schistosomosis management. But among the research that has been interested in molecules, it is the impact of treatment with praziquantel that is the most evaluated, by 50% of these studies. It is the only molecule combined with albendazole that is currently used by the NSCP. So, any subsequent investigation of the program consists of evaluating the impact of praziquantel treatment.

The literature review revealed that only thirteen (13) scientific documents (15.66%) on schistosomosis in Burkina Faso had mentioned their funding source. Our results are different from Makaula et al. [102] who found that in Malawi, 98% of the research carried out on schistosomosis were funded. In terms of funding sources, of the thirteen documents, USAID and AUF were the main ones. Although the challenge of schistosomosis in Burkina Faso was first a national public health

problem before being international, local funding was almost non-existent. This low declination of identities by sponsors who supported the research in Burkina Faso, would be justified by political and strategic reasons.

At the national level, the prevalence of schistosomosis increased from 62.4% in the 1970s to 52.86% just before the implementation of the program in 2002 and significantly decreased over the years. Toure et al. [103] in 2015 in Burkina Faso noted a considerable decrease in the national prevalence of *S. haematobium* infection. This rate was 59.6% before 2004. Two years after the implementation of the program among children from 5 to 15 years old, it decreased to 7.7%. All this shows the impact that the NSCP generates in the process of schistosomosis control and elimination in Burkina Faso.

Our study that aimed at giving a general idea of the work which was carried out on schistosomosis in Burkina Faso showed that original scientific research articles were the most frequently found types of documents over the period from 1960 to 2020. Most of the studies were conducted by Burkinabe authors. Urogenital schistosomosis and intestinal schistosomosis with their respective etiological agent attracted very much drew the authors' attention. References to sources of funding were rarely included in the documents. Overall, we noted a decrease in the prevalence of schistosomosis probably largely related to the NSCP implementation. The NSCP effectiveness could therefore contribute to the elimination of schistosomosis in Burkina Faso.

## References

- [1] Aubry P., Gaüzere A.B. 2015. Schistosomoses ou bilharzioses. *Medecine Tropicale* 8(7): 1–80.
- [2] van der Werf M.J., de Vlas S.J., Brooker S., Looman C.W., Nagelkerke N.J.D., Habbema J.D.F., Engels D. 2003. Quantification of clinical morbidity associated with schistosome infection in sub-Saharan Africa. *Acta Tropica* 86(2–3): 125–139. doi:10.1016/s0001-706x(03)00029-9
- [3] WHO. 1993. The control of schistosomiasis, second report of the WHO expert committee. Technical Report Series n° 830. <https://www.who.int/publications/i/item/WHO-TRS-830>
- [4] WHO. 2006. Weekly epidemiological record 81 (51/52): 469–480. <https://iris.who.int/bitstream/handle/10665/240837/>

- WER8151\_52.PDF
- [5] Pelligrino A., Giudicelli P. 1961. Radiodiagnostic dans la bilharziose urinaire. *Gazette Médicale de France* 68: 2155–2166.
- [6] Doumenge J.P., Mott K.E., Cheung C., Villenave D., Chapuis O., Perrin M.F., Reaud-Thomas G. 1987. Atlas of the global distribution of schistosomiasis. CEGET-CNRS-OMS-WHO. Presses Universitaires de Bordeaux, France.
- [7] Gioda A. 1992. Les mêmes causes ne produisent pas les mêmes effets: travaux hydrauliques, santé et développement. *Sécheresse* 3(4): 227–234.
- [8] Poda J.N., Sondo B., Parent G. 2003. Influence des hydro-aménagements sur la distribution des bilharzioses et de leurs hôtes intermédiaires au Burkina Faso. *Cahiers Sante* 13: 49–53.
- [9] Ministère de la Santé du Burkina Faso. 2003. Programme National de Lutte contre la schistosomiase. Rapport Technique: 1–48.
- [10] Katherine T., Fridman J.F. 2007. An update on anemia in less developed countries. *American Journal of Tropical Medicine and Hygiene* 77(1): 44–51.
- [11] OMS et UNICEF. 2004. La schistosomiase et les geohelminthiases: actions de prévention et de lutte. WHO/CDS/CPE/PVC/ 9: 1–4.
- [12] Sawadogo M. 2004. Diagnostic biologique des schistosomiasis intestinale et urogénitale: cas de la commune de Mogtedo (Province de Ganzourgou). PhD Thesis, Université de Ouagadougou, Burkina Faso. n° 892.
- [13] Vuitton D.A., Bresson H.S., Delabrousse E., Manton G.A. 2004. Foie et maladies parasitaires. *Gastroentérologie Clinique et Biologique* 28: 1122–1137.
- [14] Ministère de la Santé. 2006. Direction des études et de la planification, Burkina Faso. Annuaire statistique de santé.
- [15] Becket R., Sout J. 1969. La bilharziose intestinale à *Schistosoma intercalatum* en Haute-Volta. *Bulletin de la Société de Pathologie Exotique* 62(1): 146–151.
- [16] Boudin C., Simonkovich E. 1979. Enquête parasitologique sur les bilharzioses humaines dans la région de Banfora en Haute-Volta. *Doc. tech OCCGE* n°6777. [https://horizon.documentation.ird.fr/exl-doc/pleins\\_textes/pleins\\_textes\\_5/b\\_fdi\\_08-09/09748.pdf](https://horizon.documentation.ird.fr/exl-doc/pleins_textes/pleins_textes_5/b_fdi_08-09/09748.pdf)
- [17] Boudin C., Moreau J.P. 1979. Essai de traitement de masse de la bilharziose intestinale à *Schistosoma mansoni* par prise unique répétée d'oxamnique. *Doc. tech OCCGE* n°6841. [https://horizon.documentation.ird.fr/exl-doc/pleins\\_textes/pleins\\_textes\\_5/b\\_fdi\\_08-09/09751.pdf](https://horizon.documentation.ird.fr/exl-doc/pleins_textes/pleins_textes_5/b_fdi_08-09/09751.pdf)
- [18] Boudin C. 1979. Enquête de prévalence sur la schistosomiase urinaire dans la région du sahel voltaïque. *Doc. tech OCCGE* n°7220. [https://horizon.documentation.ird.fr/exl-doc/pleins\\_textes/pleins\\_textes\\_5/b\\_fdi\\_02-03/00257.pdf](https://horizon.documentation.ird.fr/exl-doc/pleins_textes/pleins_textes_5/b_fdi_02-03/00257.pdf)
- [19] Colette J., Sellin B., Plouvier S., Leroy J.C., Frohon J., Perrau D. 1977. Résultats de sondage parasitologique effectués chez les élèves de l'école de la vallée du kou. *ORSTOM* n°8570.
- [20] Sellin B., Simonkovich E., Diarrassouba Z. 1979. Les mollusques hôtes intermédiaires des schistosomes dans les secteurs de Dori, Kaya, Ouahigouya et Dédougou. *Doc. tech OCCGE* n°28775.
- [21] Sellin B., Desfontaine M. 1980. Recherche d'une méthode de lutte contre la schistosomiase à *Schistosoma haematobium* en zone de savane sèche Haute-Volta. *Doc. tech OCCGE* n°7299.
- [22] Sellin B., Simonkovich E., Roux J. 1980. Etude de la répartition des mollusques hôtes intermédiaires des schistosomes en Afrique de l'Ouest. Premiers résultats. *Médecine Tropicale* 40(1): 31–40. <https://core.ac.uk/download/pdf/39877833.pdf>
- [23] Boudin C., Sellin B., Simonkovich E. 1981. Enquête sur la prévalence des bilharzioses dans les régions de Kombissiri et Ziniaré, Haute-Volta en Mars 1978. *Doc. tech OCCGE* n°6778/78. [https://horizon.documentation.ird.fr/exl-doc/pleins\\_textes/pleins\\_textes\\_5/b\\_fdi\\_02-03/00238.pdf](https://horizon.documentation.ird.fr/exl-doc/pleins_textes/pleins_textes_5/b_fdi_02-03/00238.pdf)
- [24] Boudin C., Capron A., Dessaint J.P. 1981. Immunologie humorale, immunologie cellulaire, épidémiologie et thérapeutique dans les schistosomiasis. *ORSTOM fonas documentaire* n°28649 cote B.
- [25] Sellin B., Boudin C. 1981. Les schistosomiasis en Afrique de l'Ouest. *Etudes Médicales* 1: 1–89.
- [26] Belem A.P. 1982. Contribution à l'étude de la bilharziose à *S. mansoni* et essai d'un traitement de masse par Vansil en Haute-Volta. PhD Thesis, Faculty of Medicine, University of Abidjan. n°349.
- [27] Colette J., Sellin B., Garrigue G., Simonkovich E. 1982. Etude épidémiologique de la substitution de *Schistosoma haematobium* par *Schistosoma mansoni* dans une zone d'endémie bilharzienne d'Afrique de l'Ouest (Haute-Volta). *Médecine Tropicale* 42(3): 289–296.
- [28] Sellin B., Simonkovich E., Ovazza L., Sellin E., Desfontaine M., Rey J.I. 1982. Valeur de l'examen macroscopique des urines et les bandelettes réactives pour la détection de l'hématurie et de la protéinurie dans le diagnostic de masse de la schistosomiase urinaire, avant et après le traitement. *Médecine Tropicale* 42(5): 521–528.
- [29] Bresseur P., Drhuile P. 1984. Dispersion géographique de l'endémie bilharzienne dans trois départements du Burkina Faso. *Bulletin de la Société de Pathologie Exotique* 77(5): 673–677.
- [30] Le Bras M., Commenges D., Dupont A., Giap G., Guerin B., Ouedraogo J.B., Traore A., Villenave D. 1986. Étude critique d'une méthode d'enquête interdisciplinaire sur la schistosomiase urinaire au Burkina Faso. *Bulletin de la Société de Pathologie*



- Exotique* 79(3): 398–408.
- [31] Compaore H. 1987. Infestation bilharzienne a *Schistosoma haematobium* dans la province du Bam, Burkina Faso. PhD Thesis, University of Ouagadougou, Burkina Faso. n°409.
- [32] Ouiminga R.M., Diallo M., Tiendrebeogo A., Traore L.K., Soudre R.B. 1987. Bilharziose cutanée, à propos d'une observation au CHUYO, Burkina Faso. *Médecine d'Afrique Noire* 34(6): 571–572.
- [33] Mazer A., Sankale M. 1988. Guide de médecine en Afrique et Océan Indien. EDICEF.
- [34] Ouedraogo N. 1984. Les facteurs de risque de la schistosomiase urinaire dans la région de Kaya, Burkina Faso. PhD Thesis, University of Ouagadougou, Burkina Faso n°422.
- [35] Traore L.K., Ouedraogo L.H., Pietra V., Nacoulma I., Nebie B., Pafadnam D.E. Salles F. 1990. Prévalence de l'infection a *Schistosoma haematobium* et relation bilharziose hématurie dans deux villages du Burkina Faso, *Médecine d'Afrique Noire* 37(3): 100–107.  
<http://www.santetropicale.com/Resume/33704.pdf>
- [36] Schilling C., Nougara A., Korte R. 1992. Diagnostic de la bilharziose urinaire à l'aide de méthode combines de dépistage de masse, Burkina Faso, *Médecine d'Afrique Noire* 39(4): 251–256.  
<http://www.santetropicale.com/Resume/43901.pdf>
- [37] Zan S. 1992. Enquête sanitaire de base dans la zone d'aménagement hydro agricole et hydroélectrique de Bagré, Burkina Faso: a propos d'une étude sur les schistosomiasis et les autres parasitoses intestinales majeures liées à l'hygiène de l'eau. PhD Thesis, University of Ouagadougou, Burkina Faso. n°201.
- [38] World Health Organization. 1993. The control of schistosomiasis: second report of the WHO expert committee. WHO Technical Report. Series 830.
- [39] Bouree P., Cornet M., Romand S., Bisaro F. 1994. Bilharziose intestinale avec rectorragies: à propos de trois cas. *La Revue de Médecine Interne* 15(7): 481–483. doi:10.1016/S0248-8663(05)81473-0
- [40] Poda J.N., Sellin B., Sawadogo L. 1994. Dynamique des populations de *Bulinus senegalensis* dans une mare temporaire située dans une zone climatique nord soudanienne au Burkina Faso. *Revue d'élevage et de Médecine Vétérinaire des Pays Tropicaux* 47(4): 375–378.
- [41] Poda J.N., Sellin B., Sawadogo L. 1994. Hôtes intermédiaires et prévalence de la bilharziose au Burkina Faso. *Science et Technique* 20: 54–67.
- [42] Poda J.N., Sellin B., Sawadogo L., Sanogo S. 1994. Distribution spatiale des mollusques hôtes intermédiaires potentiels des schistosomes et de leurs biotopes au Burkina Faso. *OCCGE Information* 101: 12–19.  
[https://horizon.documentation.ird.fr/exl-doc/pleins\\_textes/pleins\\_textes\\_6/b\\_fdi\\_35-36/41523.pdf](https://horizon.documentation.ird.fr/exl-doc/pleins_textes/pleins_textes_6/b_fdi_35-36/41523.pdf)
- [43] Tiendrebeogo S.R. 1994. Parasitoses intestinales et bilharziose urinaire en milieu scolaire dans la ville de Ouagadougou. PhD Thesis, University of Ouagadougou, Burkina Faso. n°04.
- [44] Klutse A., Baleux B. 1996. Etude de la survie de *Bulinus truncatus* et de *B. pfeifferi* dans les eaux usées épurées par lungage en zone soudano sahéenne, Ouagadougou. *Médecine Tropicale* 56(1): 41–47.
- [45] Poda J.N. 1996. Distribution spatiale des hôtes intermédiaires des schistosomes au Burkina Faso: facteurs influençant la dynamique des populations de *Bulinus truncatus rohlfsi* et de *Bulinus senegalensis*. PhD Thesis, University of Ouagadougou, Burkina Faso.
- [46] Poda J.N., Sawadogo L.L., Sellin B., Sanogo S. 1996. Dynamique des populations de *Bulinus truncatus rohlfsi* dans le barrage de Dyono en zone nord soudanienne du Burkina Faso. *Agronomie Africaine* 8(1): 61–68.
- [47] Campagne G., Poda J.N., Garba A., Bremond P., Labbo R., Campaore I., Kambire R., Parent G., Chippaux J.P. 1998. Le risque bilharzien dans la région du barrage de Bagré, Burkina Faso. *Médecine Tropicale* 58(4): 415–416.
- [48] Elcuar R., Armas M., Ramirez M., Noguera F.J., Bolanos M., Quinones I., Lafarga B. 1998. Outbreak of schistosomiasis in a group traveler returning from Burkina Faso. *Enfermedades Infecciosas y Microbiologia Clinica* 16(8): 367–369.
- [49] Garba A., Campagne G., Poda J.N., Parant G., Karambiri R., Chippaux J.P. 1999. Les schistosomoses dans la région de Ziga au Burkina Faso avant la construction du barrage. *Santé Publique* n°2043.
- [50] Sankande B., Traore S.S., Kabore J., Ouattara T., Soudre R.B. 1998. Parasitoses humaines au Burkina Faso: approche histopathologique, cas des schistosomes. *Bulletin de la Societe de Patologie Exotique* 91(3): 217–220.
- [51] Poda J.N., Traore A. 2000. Situation des schistosomes au Burkina Faso: la lutte contre les schistosomes en Afrique de l'Ouest. *IRD Paris (FRA)*: 177–195.
- [52] Poda J.N., Daniou D., Kambou T., Sawadogo B., Et Sondo B. 2001. Etude comparative de 3 foyers bilharziennes à *Schistosoma haematobium* au Burkina Faso. *Bulletin de la Societe de Patologie Exotique* 94(1): 25–28.
- [53] Poda J.N., Sorgho H., Dianou D., Sawadogo B., Kambou T., Parent G., Sondo B. 2001. Profil parasitologique de la schistosomose urinaire du complexe hydro agricole du Sourou au Burkina Faso. *Bulletin de la Societe de Patologie Exotique* 94(1): 21–24.
- [54] Ministère de la Santé du Burkina Faso. 2003. Programme national de lutte contre la schistosomiase. Rapport Technique:1–48.
- [55] Poda J.N., Sondo B., Parent G. 2003. Influence des

- hydro-aménagements sur la distribution des bilharzioses et de leurs hôtes intermédiaires au Burkina Faso. *Médecine et Sante Tropicales* 1(13): 49–53.
- [56] Dianou D., Poda J.N., Savadogo L.G., Sorgho H., Wango S.P., Sondo B. 2004. Parasitoses intestinales dans la zone du complexe hydro-agricole du Sourou au Burkina Faso : cas de schistosomose intestinale. *La Revue Electronique en Science de l'Environnement* 5(2): 1–8. doi:10.4000/vertigo.3369
- [57] Nacoulma J.R.F. 2004. Evolution de la schistosomiase dans le district sanitaire de Tougan de 1999 à 2003. Mémoire Attaché de Santé en Epidémiologie, ENSP Ouagadougou Burkina Faso. <https://www.memoireonline.com/07/08/1221/evolution-schistosomiase-dans-le-district-sanitaire-de-tougan.html>
- [58] Poda J.N., Traore A., Sondo B.K. 2004. Endémie bilharzienne au Burkina Faso. *Bulletin de la Societe de Patologie Exotique* 97(1): 47–52.
- [59] Poda J.N., Wango S.P., Sorgho H., Dianou D. 2004. Evolution récente des schistosomoses dans le complexe hydro-agricole du Sourou au Burkina Faso. *Bulletin de la Societe de Patologie Exotique* 97(1): 15–18.
- [60] Sawadogo M. 2004. Diagnostic biologique des schistosomiasis intestinale et urogénitale et recherche des hôtes intermédiaires: cas de la commune de Mogtedo (Province de Ganzourgou). PhD Thesis, University of Ouagadougou, Burkina Faso. n° 892.
- [61] Vuitton D.A., Bresson H.S., Delabrousse E., Mantion G.A. 2004. Foie et maladies parasitaires dans cahier FMC. *Gastroentérologie Clinique et Biologique* 28: 1122–1137.
- [62] OMS. 2006. Weekly epidemiological record. <http://www.who.int/wer>. 81 (16): 145-164.
- [63] Kabore I. 2005. Evaluation microscopique de terrain pour le diagnostic des schistosomiasis et autres parasitoses intestinales. PhD Thesis, University of Ouagadougou, Burkina Faso. n°1715.
- [64] Poda J.N., Mwanga J., Daniou D., Garba A., Ouedraogo F.C., Zongo D., Sondo B.K. 2006. Les parasitoses qui minent les nouveaux pôles de développement au Burkina Faso: cas des schistosomoses et des géohelminthes dans le Sourou. *La Revue Electronique en Sciences de l'Environnement vertigo* 17(2): 1–7.
- [65] Sorgho H. 2006. Profil parasitologique de la schistosomiase et sérodiagnostic de la schistosomiase intestinale dans deux hydro-aménagement à l'ouest du Burkina Faso. PhD Thesis, University of Ouagadougou, Burkina Faso.
- [66] Koukounari A., Gabielli A.F., Toure S., Olivia E.B., Zhang Y, Sellin B., Donnelly C.A., Fenwick A., Webster J.P. 2007. *Schistosoma haematobium* infection and morbidity before and after large scale administration of praziquantel in Burkina Faso between 2004–2005. *Journal of Infectious Diseases* 196: 259–269.
- [67] Poda J.N. 2007. Les maladies liées à l'eau dans le bassin de la Haute-Volta: état des lieux et perspectives, cas de schistosomes. Volta Basin Focal Project Report (4): 87.
- [68] Poda J.N., Mwanza J., Dianou D., Garba A., Ouedraogo F.C., Zongo D., Sondo B.K. 2007. Les schistosomoses au complexe hydro agricole du Sourou au Burkina Faso: situation et modèle de transmission. *Développement, Environnement et Santé*: 1-14
- [69] Zongo D., Kabre B.G., Poda J.N., Dianou D., Sellin B. 2007. Evolution à long terme de la situation de la schistosomiase urinaire dans le village de Yamakoro au Burkina Faso. *Science et Technique, Sciences de la Santé* 30(1–2): 27–33.
- [70] Traore A., Barro-Traore F., Goumbri Lompo O., Niamba P., Diato G.N., Tapsoba P., Nikiema V., Ilboudo L, Bonkougou M., Heid E., Grossahans E., Soudre B.R. 2008. Bilharziose cutanée. Six cas. *Annales de Dermatologie et de Vénérologie* 135(1): 71–72. doi:10.1016/j.annder.2007.06.005
- [71] Gouba W.N. 2009 Evolution des marqueurs immunologiques de l'infection a *Schistosoma mansoni* chez les patients après traitement au praziquantel a la vallée du kou au Burkina Faso. Mémoire de DEA en Biologie Appliquée et Modélisation des Systèmes Biologiques. Université Polytechnique de Bobo Dioulasso, IRD/IRSS. <https://beep.ird.fr/collect/upb/index/assoc/IDR-2009-GOU-EVO/IDR-2009-GOU-EVO.pdf>
- [72] Ido K.B. 2009. Apport de l'échographie dans le bilan lésionnel et l'évaluation de la morbidité de la bilharziose urinaire chez l'enfant, à propos de 590 enfants des régions de l'Est et du centre est du Burkina Faso. PhD Thesis, University of Ouagadougou. n°111.
- [73] Zongo D., Kabre B.G., Sawadogo B., Poda J.N. 2009. Importance of malacological factor in the transmission of *Schistosoma haematobium* in two dams in the province of Oubritenga, Burkina Faso. *Journal of Environment Sciences* 3(1): 127–133.
- [74] Dicko H. 2009. Etude de la prévalence et de l'intensité de l'infection a schistosoma chez les femmes enceintes et les filles en âge de procréer dans deux régions du Burkina Faso. PhD Thesis, University of Ouagadougou, Burkina Faso. n°1637.
- [75] Toure S. 2010. Schistosomiasis in women of reproductive age in Burkina Faso; implications for control. Institute for Health Research and Development, ClinicalTrials.gov. n° NCT00463632.
- [76] Koukounari A., Toure S., Donnelly C.A., Ouedraogo A., Yoda B., Ky C., Kabore M., Bosque-Oliva E., Basanez M.G., Fenwick A., Webster J.P. 2011. Integrated monitoring and evaluation and environmental risk factors for urogenital

- schistosomiasis and active trachoma in Burkina Faso before preventive chemotherapy using sentinel sites. *BMC Infectious Diseases* 11: article number 191. doi:10.1186/1471-2334-11-191
- [77] Nana A. 2012. Évaluation de la schistosomiase chez les enfants d'âge scolaire avant la mise en eau du barrage du Soum au Burkina Faso. Mémoire de Master 2 en Parasitologie Médicale, Université Polytechnique de Bobo Dioulasso, Burkina Faso. n°004.
- [78] Ouedraogo H., Kabore A., Down P., Ottesen E. 2012. Isolated and persistent intestinal schistosomiasis foci in the western province of Burkina Faso. ASTMH, Atlanta, GA November 2012.
- [79] Siri B.A., Yonli A., Yaka P., Coulibali S. 2012. Etude de la relation des paramètres météorologique et de l'incidence des schistosomiasis urinaires dans la région du centre du Burkina Faso de 2006–2010. Santé et Mutations Environnementales dans les Pays en Développement. [https://jssb-bf.org/docs/abstract\\_books/Book\\_Abstract\\_JSSB\\_2012.pdf](https://jssb-bf.org/docs/abstract_books/Book_Abstract_JSSB_2012.pdf)
- [80] Gatta L.B., Balzarina P., Cadei M., Castelli K., Grigolato F. 2012. *Haematobium* eggs detection in human bladder cancer and sporocysts in snail vectors: seven cases report and a review of the Burkina Faso literature. *Pathologie-Biologie* 60(5): 287–290. doi:10.1016/j.patbio.2011.10.009
- [81] Zongo D., Kabré B.G., Dayeri D., Sawadogo B., Poda J.N. 2012. Etude comparative de la transmission de la schistosomiase (formes urinaire et intestinale) dans dix sites du Burkina Faso. *Médecine et Santé Tropicales* 22(3): 323–329. doi:10.1684/mst.2012.0089
- [82] Kpoda N.W., Sorgho H., Ouedraogo J.B., Kabre B.G. 2013. Endémie bilharzienne a *Schistosoma mansoni* dans la vallée du kou au Burkina Faso: caractérisation du système de transmission et impact socio-économique. *Comptes Rendus Biologie* 336(5): 284–288. doi:10.1016/j.crv.2013.04.008
- [83] Traore I. 2013. geographic determinants of human schistosomiasis transmission in the Sourou valley, Burkina Faso. PhD Thesis, Georg-August University, Göttingen <http://hdl.handle.net/11858/00-1735-0000-0001-BB25-F>
- [84] Bagayan M., Zongo D., Savadogo B., Ouedat A., Sorgho H., Kabre B.G., Poda J.N. 2014. Prévalence de la schistosomiase urinaire en milieu scolaire dans un milieu péri urbain de Ouagadougou: cas du quartier Yamtenga (Burkina Faso). *Science et Technique, Sciences de la Santé* 37(1–2): 93–100.
- [85] Bugge H.F. 2014. Control and elimination strategies for schistosomiasis in Burkina Faso, West Africa. The effect of five years of mass drug administration on prevalence and intensity of infection. Student Thesis, Faculty of Medicine, University of Oslo. [https://www.duo.uio.no/bitstream/handle/10852/43422/1/Studentoppagave-Bugge-Hovmork\\_031014\\_Innlevert.pdf](https://www.duo.uio.no/bitstream/handle/10852/43422/1/Studentoppagave-Bugge-Hovmork_031014_Innlevert.pdf)
- [86] Hafiz L. 2014. Geostatistical modelling for the monitoring of environmental related endemic disease: applicaton to the distribution of schistosomiasis in Burkina Faso. Master Thesis, École Polytechnique Fédérale de Lausanne. <https://infoscience.epfl.ch/record/203586>
- [87] Napon A.M., Bandre E., Nikiema Z., Diallo O., Ouedraogo I., Kaboret S.D., Lougue Sorgho L.C., Alfidja A., Boyer L. 2014. Child abdominal tumor in tropical context: think about schistosomiasis. *African Journal of Paediatric Surgery* 11(2): 194–195. doi:10.4103/0189-6725.132838
- [88] Perez-Saez J., Ceperley N., Mande T., Frhlich J.M., Rinaldo A. 2014. Comprendre le mode de transmission de la bilharziose au Burkina Faso. Ecole Polytechnique Fédérale de Lausanne. <https://www.epfl.ch/labs/echo/page-12894-fr-html/labs-echo-fr-page-12894-fr-html-page-12895-fr-html-page-113944-fr-html/>
- [89] Savadogo L.B.G., Savadogo B., Dianou D., Poda J.N. 2014. *Schistosoma mansoni* and *Schistosoma haematobium* infection and nutritional status of children in the hydro-agricol complex of Sourou. *Occupational Diseases and Environmental Medicine* 2(2): 13–18. doi:10.4236/odem.2014.22002
- [90] Zongo D., Kabre B.G., Poda J.N., Dianou D. 2014. Schistosomiasis among farmer and fisherman in the west part of Burkina Faso. *Journal of Biology Sciences* 8(2): 482–485. doi:10.3923/jbs.2008.482.485
- [91] Zongo D., Bagana M., Sawadogo B., Drabo F., Ouedraogo A., Poda J.N. 2015. Endémie de la schistosomiase et des géohelminthes chez les enfants au Burkina Faso. *Sciences de la Santé* 38(1–2). [https://revuesciences-techniquesburkina.org/index.php/sciences\\_de\\_la\\_sante/article/view/636](https://revuesciences-techniquesburkina.org/index.php/sciences_de_la_sante/article/view/636).
- [92] Perez-Saez J., Mari L., Bertuzzo E., Casagrandi R., Gatto M., Rinaldo A., Sokolow S.N., Deleo G.A., Mande T., Fraochlich J.M., Karambiri H., Yacouba H., Maiga A., Sou M. 2015. A theoretical analysis of the geography of schistosomiasis in Burkina Faso high lights the roles of human mobility and water resources development in disease transmission. *PLOS Neglected Tropical Diseases* 9(10): e0004127. doi:10.1371/journal.pntd.0004127
- [93] Diallo A.C. 2015. Etude de l'impact du traitement de masse sur l'évolution des schistosomoses au Burkina Faso de 2004 à 2013. PhD Thesis, University of Ouagadougou, Burkina Faso.
- [94] Faes A. 2015. Effect of water contacts seasonality on schistosomiasis transmission in rural sahelian area in Burkina Faso. Master Thesis, École Polytechnique Fédérale de Lausanne.

- <https://infoscience.epfl.ch/record/213509>
- [95] Perez-Saez J., Bertuzzo E., Frogelich J.M., Mande T., Ceperley M., Sou M., Yacouba H., Maiga H., Sokolow S., Casagrangi R., Gatto M., Mari L., Rinaldo A. 2015. Spatial patters of schistosomiasis in Burkina Faso: relevance of human mobility resources development. *Geophysical Research Abstracts* Vol. 17, EGU2015-6328, 2015 EGU General Assembly 2015.
- [96] Savadogo B., Adjami A.G., Zongo D., Hattendorf J., Poda J.N., Traore A.S. 2015. Schistosomiasis assessment among school children and macological factors in Burkina Faso: case of Daguilma, Koubri, Yamtenga. *Current Research in Microbiology and Biotechnology* 3(4): 701–706.
- [97] Ministère de la santé. 2016. Plan stratégique de lutte contre les maladies tropicales négligées 2016–2020. [https://espen.afro.who.int/system/files/content/resources/SENEGAL\\_NTD\\_Master\\_Plan\\_2016\\_2020.pdf](https://espen.afro.who.int/system/files/content/resources/SENEGAL_NTD_Master_Plan_2016_2020.pdf).
- [98] WHO. 2016. Schistosomiasis et géohelminthiase: nombre de personne traitées en 2016. [http://www.who.int/schistosomiasis/genital\\_schistosomiasis/en/](http://www.who.int/schistosomiasis/genital_schistosomiasis/en/).
- [99] Ouedraogo H., Drabo F., Zongo D., Bagagna M., Bamba I., Pima T., Wienne F.Y., Toubali E., Zhang Y. 2016. Schistosomiasis in school age in Burkina Faso a decade of prevention chemotherapy 2004–2013. *Bulletin of the World Health Organization* 94(1): 37–45. doi:10.2471/BLT.15.161885
- [100] Traore I., Karthe D., Sie A., Coulibaly V., Kappas M. 2016. Influence of air temperature of children water contacts with respect to schistosomiasis transmission, risk in the Sourou valley, Burkina Faso. *Stochastic Environmental Research and Risk Assessment* 30: 2097–2107. doi:10.1007/s00477-016-1214-x
- [101] Zida A., Briegel J., Wacker J., Kabre I., Sawadogo M.P., Sanganre I., Bamba S., Yacouba A., Ouedraogo A., Yonli D., Drabo F., Traore L.K., Ouedraogo R.T., Guiguemde T.R. 2016. Epidemiological and clinical aspect of urogenital schistosomiasis in the women of Burkina Faso, west Africa. *Infectious Diseases of Poverty* 5: article number 81. doi:10.1186/s40249-016-0174-1
- [102] Makaula P., Sadalaki J.R., Muula A.S., Kayuni S., Jemu S., Bloch P. 2014. Schistosomiasis in Malawi: a systematic review. *Parasites and Vectors* 7: article number 570. doi:10.1186/s13071-014-0570-y
- [103] Toure S., Zhang Y., Bosque-Oliva E., Ky C., Ouedraogo A., Koukounari A., Gabriella A.F., Sellin B., Webster J.P., Fenwick A. 2008. Two-year impact of single praziquantel treatment on infection in the national control programme on schistosomiasis in Burkina Faso *Bulletin of the World Health Organization* 86(10): 737–816. doi:10.2471/BLT.07.048694

Received 01 November 2023

Accepted 05 February 2024